Aero 320: Numerical Methods

Homework 3

Name:	
-------	--

Due: October 7, 2013

NOTE: All problems, except the last, are to be done by hand (with the help of a calculator) but you need to show all the steps. Turn in a hard copy of your HW stapled with this as cover sheet with your name written in the above field. Submit your HW by Monday midnight at Room 201, Reed McDonald Building. Late submissions or failure to submit in the required format will receive no credit.

Problem 1

Basics on matrix algebra

 $(3 \times 7 = 21 \text{ points})$

Find whether the following statements are *true* or *false*. If you think a statement is *true*, then prove that statement. If *false*, then provide a counterexample.

- (a) Any matrix can be written as the sum of a symmetric and a skew-symmetric matrix.
- (b) Let the square matrices M and N are of same size. If $\det(M) = 0$, and $\det(N) = 0$, then $\det(M+N) = 0$.
- (c) Given square matrices X and Y of same size, $\operatorname{tr}(XY-YX)$ need not be zero.

(d) Let
$$A = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$
, and $B = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$. A and B have same eigenvalues.

- (e) Consider the matrices in part (d). Eigenvalues of AB are the *product* of eigenvalues of A and eigenvalues of B. Eigenvalues of A + B are the sum of eigenvalues of A and eigenvalues of B.
- (f) Determinant of any orthogonal matrix is ± 1 .

(g) Let
$$A = \begin{pmatrix} 0 & 1 & 2 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix}$$
. Then A^{2013} is $\begin{pmatrix} 0 & 0 & 0 \\ 1 & 1 & 1 \\ 2 & 2 & 2 \end{pmatrix}$.

Problem 2

Uniqueness of the solution for a system of linear equations $(3 \times 3 = 9 \text{ points})$

Show that the following system of equations:

$$x_1$$
 + $4x_2$ + αx_3 = 6,
 $2x_1$ - x_2 + $2\alpha x_3$ = 3,
 αx_1 + $3x_2$ + x_3 = 5,

has (a) unique solution if $\alpha = 0$, (b) no solution when $\alpha = -1$, and (c) infinitely many solutions when $\alpha = 1$.

Problem 3

Gauss elimination for an ill-conditioned system (5+5+5=15 points)

Consider a system of linear equations that is "ill-conditioned", meaning it may be difficult to obtain the numerical solution accurately (see Section 2.4 in textbook). One such system, written in augmented matrix form, is

$$\begin{pmatrix}
3 & 2 & 4 & 9 \\
8 & -6 & -8 & -6 \\
-1 & 2 & 3 & 4
\end{pmatrix}.$$

- (a) By doing Gauss elimination using exact arithmetic (use fractions throughout), show that the exact solution is $x = \{1, 1, 1\}^{\top}$.
- (b) Now compute the solution using Gauss elimination with only 3 significant digits in each operation. How does your answer compare with part (a)?
- (c) Compute the solution to the system by changing the coefficient in row 1, column 1 position to 3.1 instead of 3. Use your calculator's full precision. How does this answer compare to the previous solutions in part (a) and (b)?

Problem 4

Gauss elimination algorithm: standard versus partial pivoting (5+2+3+5=15 points)

Let

$$A = \begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{pmatrix}, \qquad b = \begin{pmatrix} 1 \\ 3 \\ 5 \end{pmatrix}.$$

- (a) Show that the system of linear equations Ax = b has infinitely many solutions.
- (b) Describe the set of all possible solutions.
- (c) Use Gauss elimination to solve Ax = b using exact arithmetic. Because there are infinitely many solutions, it is unreasonable to expect one particular solution to be computed. What does happen?
- (d) Repeat part (c) with partial pivoting. What are your conclusions?

Problem 5

Forces along the links of a truss

(10 + 30 = 40 points)

The following is a schematic of a mechanical structure (such as a bridge), modeled by the truss as follows.




Figure 1: Schematic of the truss structure.

As shown in the diagram, the truss has 10 joints, numbered in circles. It has 17 rigid links, also numbered in the diagram along the line segments. Given the loads F_1, F_2, F_3, F_4 , our objective is to compute the forces f_1, f_2, \ldots, f_{17} along the rigid links of the structure.

To do this, let us introduce $\alpha = \sin(45^{\circ}) = \cos(45^{\circ})$. Now, for each joint, we write the force balance equations in horizontal and vertical directions:

Joint 2
$$\begin{cases} -\alpha f_1 + f_4 + \alpha f_5 = 0, \\ -\alpha f_1 + f_3 + \alpha f_5 = 0, \end{cases}$$
Joint 3
$$\begin{cases} -f_2 + f_6 = 0, \\ -f_3 + F_1 = 0, \end{cases}$$
Joint 4
$$\begin{cases} -f_4 + f_8 = 0, \\ f_7 = 0, \end{cases}$$
Joint 5
$$\begin{cases} -\alpha f_5 - f_6 + \alpha f_9 + f_{10} = 0, \\ -\alpha f_5 - f_7 + \alpha f_9 + F_2 = 0, \end{cases}$$
Joint 6
$$\begin{cases} -f_8 - \alpha f_9 + f_{12} + \alpha f_{13} = 0, \\ -\alpha f_9 + f_{11} + \alpha f_{13} = 0, \end{cases}$$
Joint 7
$$\begin{cases} -f_{10} + f_{14} = 0, \\ -f_{11} + F_3 = 0, \end{cases}$$
Joint 8
$$\begin{cases} -f_{12} + \alpha f_{16} = 0, \\ f_{15} - \alpha f_{16} = 0, \end{cases}$$
Joint 9
$$\begin{cases} -\alpha f_{13} - f_{14} + f_{17} = 0, \\ -\alpha f_{13} - f_{15} + F_4 = 0, \end{cases}$$
Joint 10
$$\begin{cases} -\alpha f_{16} - f_{17} = 0. \end{cases}$$

- (a) Write the above system in the matrix equation form Ax = b, where $x = \{f_1, f_2, \dots, f_{17}\}^{\top}$.
- (b) Write a computer code to solve for x using Gauss elimination, when the external loads are
- (i) $F_1 = 10, F_2 = 15, F_3 = 0, F_4 = 10$; and (ii) $F_1 = 10, F_2 = 0, F_3 = 20, F_4 = 0$. Compare the results obtained with and without partial pivoting.